Title: Mr:	Patie	nt Addre	ec.		
			.55.		
First Name:		NHS No. (if known):			
Last Name:		GP Name and Address:			
Tolophono					
Telephone:	GP T	GP Telephone (if known):			
Gender: <b>Female</b> . D.O.B:// Age:	Would	d you lik	e us to send a copy of this consultation to your GP? $\hfill\Box$		
Patient's personal details					
Tick which of the following applies to you	Yes	No	Reconfirm details at each appointment.		
Do you have any recent or past medical history of note?					
Do you take any medicines? Antacids? Contraceptives?					
Are you currently taking any other medicines? (including any herbal remedies such as St.Johns Wort)					
Do you suffer from Bowel disease (e.g. Crohn's disease) or liver problems?					
Do you currently suffer from vomiting or diarrhoea?					
Have you ever had a serious reaction to ulipristal acetate (ellaOne) or levonorgestrel (Levonelle)?					
Sexual History					
Tick which of the following applies to you	Yes	No	Reconfirm details at each appointment.		
Have you had unprotected sex within the last 120 hours (5 days)?					
Have you had unprotected sex within the last 72 hours (3 days)?					
Have you had unprotected sex earlier in this menstrual cycle?					
Is there a possibility you may be pregnant?					
Your last menstrual period		1			
Tick which of the following applies to you	Yes	No	Reconfirm details at each appointment.		
Was your last period late, longer/shorter or unusual in any way?					
Have you already taken Levonelle or ellaOne since your last period?					
Further information about ulipristal acetate (ellaOne)		ı			
Tick which of the following applies to you	Yes	No	Reconfirm details at each appointment.		
Do you understand that if you vomit within 3 hours, another dose is required? You will need to come back or visit your doctor.					
Do you understand that If your next period is >3 days late or different in any way you should visit your doctor?					
Unprotected sex can lead to sexually transmitted diseases (STIs) do would you like further counselling?					
Write below any further information which may be r	elevant	e.g. 1	medicines taking, conditions, concerns		
Confidential sexual health patient helplines					
FPA (formerly the Family Planning Association) national he Sexual Health Line – <b>0800 567 123</b>	pline –	0845 3	3101334		

Date: \_ \_ / \_ \_ / 2 0 \_ \_

**Emergency Hormonal Contraceptive | PGD Risk Assessment Form** 

Brook Clinic – 0800 0815023 or www.brook.org.uk
Sexwise – 0800 282930 or http://www.maketherightdecision.co.uk

If you vomit within 3 hours of taking the tablet, the EHC may not have worked, you will need to return or talk to your doctor.

## **For Offical Use**

	:: [Levonelle]	or [ella	aOne] delete as appi	ropriate				
Date	Possible Pr		Referral Required?	Directions	Pharmacist	Signature		
	☐ Yes ☐ No		☐ Yes ☐ No If yes, see below.	Take immediately	/			
Reason	for referral:				<u>.</u>			
Referral	to a suitably q	ualified c	linician is required.					
Missad	Pill Advice							
Combine		If the	ree or more 30–35 microg	ram ethinylestradiol o	or two or more 20 mig	crogram		
	otive pills	ethii	If three or more 30–35 microgram ethinylestradiol or two or more 20 microgram ethinylestradiol pills have been missed in the first week of pill-taking (i.e. days 1–7) and UPSI occurred in Week 1 or the pill-free week.					
Progesto	gen-only pills	(>12	If one or more progestogen-only pills (POPs) have been missed or taken >3 hours late (>12 hours late for Cerazette®) and UPSI has occurred before a further two pills have been taken appropriately.					
ntrauteri		If co	If complete or partial expulsion is identified or mid-cycle removal of an IUD/IUS is					
contrace <sub>l</sub> Progesto	gen-only		deemed necessary and UPSI has occurred in the last 7 days.  If the contraceptive injection is late (>14 weeks from the previous injection for					
njectable		med	medroxyprogesterone acetate or >10 weeks for norethisterone enantate) and UPSI has occurred within the last 120 hours.					
Fransder contrace <sub>l</sub> Evra® Pa	otion -	prior pato	More than 2 days late starting first patch of new pack and has UPSI in week one or the prior patch free week. 7 days late starting third patch of pack, no EHC needed but om patch free week. If more than 9 days late starting second/third patch, count as UPSI. Up to 7 days late starting second patch, no EHC needed.					
	nal Advice		Davida Oralas at la		0 1 15 105 - 15 - 15 - 15			
STIs Efficacy			Barrier Contraceptives  Oral Contraceptives		Sexual health helpling IUD discussed	nes 🗆		
TIENT CO			enefits of the medicines recomm	ended and fully understan	d them. I have also had th	e apportunity to a		
			cines being given at each appoin					
ent	Name / s	signature	J	1	Date			
	for our pharmacy a	ınd/or our aı	uthorising medical agency to cor	tact you regarding custom	er satisfaction? Yes / No	0		
you consent								